

ORDERING OFFICE, ALSO FAX:

- Most recent labs
- Supporting clinicals / Recent H&P
- Insurance card, front and back

Belimumab (Benlysta)

Provider Order Form



Date:	Patient Name:	DOB:
ICD-10 code (required):		
ICD-10 description:		
<input type="checkbox"/> NKDA Allergies:		Weight lbs/kg:
Ordering Provider:		Provider NPI:
Referring Practice Name:		Phone: Fax:
Practice Address:		City: State: Zip Code:
NURSING		
<input checked="" type="checkbox"/> Provide nursing care per IVX Standard Nursing Procedures, including reaction management and post-infusion observation		
LABORATORY ORDERS		
<input type="checkbox"/> CBC	<input type="checkbox"/> at each dose	<input type="checkbox"/> every _____
<input type="checkbox"/> CMP	<input type="checkbox"/> at each dose	<input type="checkbox"/> every _____
<input type="checkbox"/> CRP	<input type="checkbox"/> at each dose	<input type="checkbox"/> every _____
<input type="checkbox"/> Other:		
PRE-MEDICATION ORDERS (ADMINISTER 30 MINUTES PRIOR TO PROCEDURE)		
<input type="checkbox"/> acetaminophen (Tylenol) <input type="checkbox"/> 500mg / <input type="checkbox"/> 650mg / <input type="checkbox"/> 1000mg PO		
<input type="checkbox"/> cetirizine (Zyrtec) 10mg PO		
<input type="checkbox"/> loratadine (Claritin) 10mg PO		
<input type="checkbox"/> diphenhydramine (Benadryl) <input type="checkbox"/> 25mg / <input type="checkbox"/> 50mg <input type="checkbox"/> PO / <input type="checkbox"/> IV		
<input type="checkbox"/> methylprednisolone (Solu-Medrol) <input type="checkbox"/> 40mg / <input type="checkbox"/> 125mg IV		
<input type="checkbox"/> Other:		
Dose:	Route:	Frequency:
INFUSION THERAPY		
<input type="checkbox"/> belimumab (Benlysta) in 250ml 0.9% sodium chloride, intravenous infusion over one hour		
▪ Dose: 10mg/kg = _____mg		
▪ Route: <input checked="" type="checkbox"/> intravenous		
▪ Frequency: <input type="checkbox"/> induction: week 0, 2, 4, and then every 4 weeks / <input type="checkbox"/> maintenance: every 4 weeks / <input type="checkbox"/> other: _____		
▪ Infuse over one hour		
<input checked="" type="checkbox"/> Flush with 0.9% sodium chloride at the completion of infusion		
<input type="checkbox"/> belimumab (Benlysta) injection		
▪ Dose: <input type="checkbox"/> 200mg		
▪ Frequency: <input type="checkbox"/> once weekly		
▪ Route: <input checked="" type="checkbox"/> subcutaneous		
<input type="checkbox"/> Patient is required to stay for 30-minute observation post infusion		
<input type="checkbox"/> Patient is NOT required to stay for observation time		
<input type="checkbox"/> Refills: <input type="checkbox"/> Zero / <input type="checkbox"/> for 12 months / <input type="checkbox"/> _____ (if not indicated order will expire one year from date signed)		
GENERAL PLAN COMMUNICATION		
Special instructions/notes:		

Ordering Provider: Initial here _____ and proceed to the next page.

ADULT REACTION MANAGEMENT

- Observe for **hypersensitivity reaction:** Fever, chills, rigors, pruritus, rash, cough, sneezing, throat irritation
- If reaction occurs:
 - Stop infusion
 - Maintain/establish vascular access
 - Notify referring provider
 - Consider giving the following PRN
 1. Acetaminophen (Tylenol) 650mg PO **OR** _____mg for pain or fever > 38 C/100.4 F
 2. Diphenhydramine (Benadryl) 25-50mg in 10ml NS slow IV push for rash, itching, pruritis
 3. Ranitidine 25mg in 10ml NS slow IV push over 5 minutes (Consider if patient already given IV Benadryl)
 4. Ondansetron (Zofran) 4mg Slow IV push over 5 minutes for nausea or vomiting.
 5. Methylprednisolone (Solumedrol) 125mg **OR** _____mg slow IV push.
 6. Other _____
 - When symptoms resolve resume infusion at 50% previous rate and increase per manufacturer's guidelines
- Severe allergic/anaphylactic reaction:**
 - If symptoms are rapidly progressing or continuing after administration of prn medications above and signs symptoms of severe allergic/anaphylactic reaction (angioedema, swelling of the mouth, tongue, lips, or airway, dyspnea, bronchospasm with or without hypotension or hypertension.)
 1. Call 911
 2. Consider giving epinephrine (1:1000 strength) 0.3ml IM. May repeat every 5-15 minutes to a maximum of 3 doses.
 3. Treat hypotension with 500ml 0.9% sodium chloride bolus. Repeat as needed to maintain systolic BP >90.
 4. Have oxygen by nasal canula available and administer 2-15 liters, titrate to keep Spo2 >92%
 5. Have Automated External Defibrillator available
 6. Notify referring provider. If unable to reach referring provider, notify Local Medical Director.
 7. Discontinue treatment

*Consider administering premedication for prophylaxis against infusion reactions and hypersensitivity reactions.

Patient Name

Patient Date of Birth

Provider Name (Print)

Provider Signature

Date

Designate the desired location and fax the order form to:

CALIFORNIA fax: (844) 889-0275 | ___ San Mateo ___ Fremont ___ San Ramon ___ TBD

ILLINOIS fax: (312) 253-7244 | ___ Glenview ___ Schaumburg ___ TBD

KANSAS / MISSOURI fax: (844) 900-1292 | ___ Overland Park, KS ___ Lee's Summit, MO ___ Briarcliff, MO ___ TBD

OHIO fax: (844) 627-2675 | ___ Dublin ___ Pickerington ___ TBD

PENNSYLVANIA fax: (844) 820-9641 | ___ Malvern ___ Bensalem ___ Montgomeryville ___ East Shore ___ West Shore ___ TBD

TENNESSEE fax: (844) 627-2518 | ___ Brentwood ___ Hendersonville ___ TBD

